

Cal/Ecotox

Exposure Factors for Bushtit (*Psaltiriparus minumus*)*

Page 1

Endpoint Type	Endpoint Value	Error	Range	Units	Sex	Life Stage	Location	Note	Reference
Body Weight - Mean	5.5	0.2 SE		g	NR	Adult	Lab	a	1
Body Weight - Mean			0.65-0.72	g	NR	Hatchling	Santa Clara; CA	b	2
Body Weight - Mean	5.37			g	NR	NR	CA	c	3
Duration of Incubation or Gestation	13			d	NR	Embryo	Santa Clara; CA	d	2
Food Ingestion Rate	35.17	1.26 SE		kJ/d	NR	Adult	Lab	e	1
Foraging Distance	See figure				B	NR	Santa Barbara; CA	f	4
Metabolic Rate	metabolic rate = 0.36(ambient T) + 14.64				NR	Adult	Lab	g	1
Metabolic Rate	3.96			cc O^2/g/h	NR	Adult	Lab	h	1
Population Density	2.2			birds/36 ha	M	Adult	Marin; CA	i	5
Population Density	71.2	49.6-92.8 95% CI		birds/40 ha	NR	NR	Madera; CA	j	6
Population Density	72.8	50.1-95.5 95% CI		birds/40 ha	NR	NR	Madera; CA	k	6
Territory Size			50-400	yd	B	Adult	Santa Clara; CA	l	2
Territory Size	0.4	0.08	0.28-0.52	ha	NR	Adult	San Mateo; CA	m	3
Time of Fledging or Metamorphosis	14			d	NR	NR	Santa Clara; CA	n	2
Time of Mating/ Laying	Mar-Apr				B	Adult	Santa Clara; CA	o	2

Notes

- a mean body weight; N=6 birds; Nov, Jun; captured at Occidental College, Los Angeles
- b range of individual body weights; N=2 birds; Stanford University Campus; See citation for body weights on days 2-7 of age.
- c mean body weight of museum specimens; N=23 birds
- d incubation period from start to hatching; N=1 clutch; Mar, Apr; Stanford University Campus
- e consumption of mealworms at 20 deg C; N=6 birds; Nov, Jun; captured at Occidental College, Los Angeles; Consumption was equivalent to 80% of bird body mass per day.
- f figure showing mapping of approximate ranges of two flocks during one season; N=2 flocks; Condition=nonbreeding; University of California campus
- g linear regression equation for relationship between nocturnal metabolism and ambient temperature; N=12 birds; Nov, Jun; captured at Occidental College, Los Angeles
- h approximate basal rate of nocturnal oxygen consumption, measured at 30 deg C; N=12 birds; Nov, Jun; captured at Occidental College, Los Angeles; Predicted standard resting metabolism for a 5.5 g bird was 4.14 cc O^2/g/h.
- i breeding density estimate; N=6 territories; Condition=breeding; Mar-May; Palomarin Field Station, Point Reyes Bird Observatory
- j density estimate on a grazed plot by transect counts using an exponential polynomial model; N=64 birds; April-May; San Joaquin Experimental Range; See citation for density estimates of other species and comparisons of other methods of abundance estimation.
- k density estimate on an ungrazed plot by point count using an exponential polynomial model; N=43 birds; April-May; San Joaquin Experimental Range
- l approximate range of nesting territory diameters; N=NR; Mar, Apr; Stanford University Campus
- m average territory size based on > 400 point observations; N=12 territories; Condition=breeding; May, Apr; Jasper Ridge Biological Experimental Area, nr Portola Valley
- n age at which young left the nest; N=1 brood; Stanford University Campus; Parental feeding continued fro 8-14 d post-fledging.
- o time of peak nesting activity; N=NR; Mar, Apr; Stanford University Campus

References

- 1 Chaplin, Susan B. 1982. The energetic significance of huddling behavior in common bushtits (*Psaltiriparus minimus*). Auk. 99:424-430.
- 2 Addicott, Alice B. 1938. Behavior of the bush-tit in the breeding season. Condor. 40:49-63.
- 3 Hertz, Paul E., J.V. Remsen, Jr. and Stacey I. Zones. 1976. Ecological and complementarity of three sympatric parids in a California oak woodland. Condor. 78:307-316.
- 4 Ervin, Stephen. 1977. Flock size, composition, and behavior in a population of bushtits (*Psaltiriparus minimus*). Bird-Band. 48(2):97-191.
- 5 DeSante, David F. 1981. A field test of the variable circular-plot censusing technique in a California coastal scrub breeding bird community, In: C. John Ralph and J. Michael Scott , eds, eds. Estimating Numbers of Terrestrial Birds; October 26-31, 1980; Asilomar, CA. Lawrence, KS: Cooper Ornithological Society; Studies in Avian Biology No. 6. p 177-185.
- 6 Verner, Jared and Lyman V. Ritter. 1985. A comparison of transects and point counts in oak-pine woodlands of California. Condor. 87:47-68.

*Cal/EPA, OEHHA and the University of California Regents are not responsible for damages of any kind resulting from the use of or reliance on information in this report. Users are encouraged to consult the original data. Updated: February 1999.